

ROME1/1422/06
 IFIC/06-04
 RM3-TH/06-16
 IFUM-862/FT

Two-loop electroweak corrections to Higgs production in proton-proton collisions

U. Aglietti ^{*},

*Dipartimento di Fisica, Università di Roma “La Sapienza” and INFN, Sezione di Roma,
P.le Aldo Moro 2, I-00185 Rome, Italy*

R. Bonciani [†],

*Departament de Fisica Teorica IFIC, CSIC, Universitat de Valencia
Edifici d’Instituts de Paterna, Apt. Correus 22085, E-46071 Valencia, Spain*

G. Degrandi [‡],

*Dipartimento di Fisica, Università di Roma Tre and INFN, Sezione di Roma III,
Via della Vasca Navale 84, I-00146 Rome, Italy*

A. Vicini [§]

*Dipartimento di Fisica, Università degli Studi di Milano and INFN, Sezione di Milano,
Via Celoria 16, I-20133 Milano, Italy*

Abstract

We study the impact of the two-loop electroweak corrections on the production of a Higgs boson via gluon-fusion in proton-proton collisions at LHC energies. We discuss the prescription to include the corrections to the hard scattering matrix element in the calculation of the hadronic cross-section $\sigma(p + p \rightarrow H + X)$. Under the hypothesis of factorization of the electroweak corrections with respect to the dominant soft and collinear QCD radiation, we observe an increase of the total cross-section from 4 to 8%, for $m_H \leq 160$ GeV. This increase is comparable with the present QCD uncertainties originating from hard scattering matrix elements.

^{*}Email: Ugo.Aglietti@roma1.infn.it

[†]Email: Roberto.Bonciani@ific.uv.es

[‡]Email: degrassi@fis.uniroma3.it

[§]Email: Alessandro.Vicini@mi.infn.it

1 Introduction

The Higgs boson is one of the missing ingredients of the Standard Model and its discovery represents one of the most important physics goals of the LHC. This goal will be achieved only if we can predict with high accuracy all the production cross sections of this particle and if we understand in detail the different decay channels and the relative backgrounds.

At the LHC, the gluon-fusion is the dominant production mode over the entire range of interesting values of the mass of the Higgs particle ($100\text{GeV} \lesssim m_H \lesssim 1\text{TeV}$). In particular, in the range $100\text{GeV} \lesssim m_H \lesssim 2m_t$ this production mode is larger by almost one order of magnitude with respect to the next important mechanism, the vector boson fusion. It is, therefore, very important to have a precise prediction of its cross section and a reliable estimate of the remaining theoretical accuracy.

The total cross section for the Higgs boson production by gluon fusion in the LO approximation was calculated in the late seventies [1]. It is an $\mathcal{O}(\alpha_s^2 \alpha)$ calculation, since the Higgs couples to the gluons only via a heavy-quark loop (the most important contribution is the one due to the loop of top). For what concerns the higher orders, the calculation of the NLO QCD corrections have been done in the infinite m_t (mass of the top) limit in [2], and, with the full quark mass dependence, in [3]. Besides of the fact that the infinite m_t approximation should be valid in the Higgs mass range $m_H \lesssim 300\text{ GeV}$, it has been noticed [4] that this approximation works also for values of m_H beyond the top threshold, and up to masses of $\mathcal{O}(1\text{TeV})$. The total effect of the NLO QCD corrections is the increase of the LO cross section by a factor 1.5–1.7, giving a residual renormalization/factorization scale dependence of about 30%. The electroweak corrections were evaluated in the infinite m_t limit in [5] and turned out to amount to less than 1%. The attention was driven by the evaluation of the NNLO QCD corrections, carried out in the infinite m_t limit by several groups [6]. The calculation shows a good convergence of the perturbative series: while the NNLO corrections are sizable, they are, nevertheless, smaller than the NLO ones. Moreover, the NNLO corrections improve the stability against renormalization/factorization scale variations. The effect due to the resummation of soft-gluon radiation has been included in [7], and the remaining theoretical uncertainty, due to higher-order QCD corrections, has been estimated to be smaller than 10%. Finally, several efforts were also devoted to the calculation of QCD radiative corrections to less inclusive quantities, such as the rapidity distribution, recently evaluated at the NNLO [8], or the transverse momentum (q_T) distribution [9], which, in [10], is evaluated using the fixed-order perturbative results up to NLO in QCD and the resummation up to the NNLL.

Motivated by this accurate scenario, the NLO electroweak corrections to the gluon fusion were reexamined recently. In [11] the contribution to the partonic cross section due to the light fermions were calculated. It turned out that they are sizeable. In particular, in the intermediate Higgs mass range, from 114 GeV up to the $2m_W$ threshold, these corrections increase the LO partonic cross section by an amount of 4–9%. For larger values of the mass of the Higgs, $m_H > 2m_W$, they change sign and reduce the LO cross section; however, in this region the light-fermion corrections are quite small, reaching at most a -2% . In [12], also the remaining electroweak corrections due to the top quark were calculated as a Taylor expansion in $m_H^2/(4m_W^2)$. They are valid for $m_H \lesssim 2m_W$, range in which they have opposite sign with respect to the light-fermion corrections. However, the corrections due




Figure 1: Lowest order (a) and generic NLO-EW (b), (c), (d) Feynman diagrams. The solid lines are fermions. The wavy lines are gauge bosons ($V = W, Z$).

to the top quark are smaller in size, reaching at most a 15% of the light-quark ones.

The impact of the NLO electroweak corrections on the hadronic cross section has not been discussed yet. We present here the effect of their inclusion in the calculation at the hadronic level.

2 Inclusion of the Two-Loop Electroweak Corrections

The partonic gluon fusion process occurs, in lowest order, via one-loop diagrams, as the one depicted in Fig. 1 (a); in the loop run only the top and the bottom quarks, because of the Yukawa suppression of the lighter quarks. The NLO-EW corrections are schematically represented by the diagrams in Figs. 1 (b), (c) and (d). In particular, in Figs. 1 (b) and (c) the WWH/ZZH couplings avoid the Yukawa suppression, and, therefore, in these diagrams the fermionic line represents all the possible flavours: light flavours, evaluated in [11], and top quark, evaluated in [12]. In Fig. 1 (d), instead, the fermionic line can represent only the top quark [12].

At the hadronic level, we consider the Higgs boson production at the LHC, and therefore in proton-proton collisions. The hadronic cross section can be written as:

$$\begin{aligned} \sigma(p + p \rightarrow H + X) &= \sum_{a,b} \int_0^1 dx_1 dx_2 f_{a,p}(x_1, M^2) f_{b,p}(x_2, M^2) \times \\ &\quad \times \int_0^1 dz \delta\left(z - \frac{\tau_H}{x_1 x_2}\right) \left(1 + \delta_{EW}(m_H)\right) \hat{\sigma}_{ab}(z) \\ \hat{\sigma}_{ab}(z) &= \hat{\sigma}_0 \left(1 + K_{ab}^{QCD \text{ only}}(\alpha_s(\mu^2), \mu^2, M^2)\right) \end{aligned} \quad (1)$$

where the partonic processes initiated by partons (a, b) are convoluted with the corresponding parton densities $f_{i,p}(x, M^2)$, ($i = a, b$), evaluated at a scale M . The effect of the higher order QCD and EW corrections is described by the two functions $K^{QCD-only}$ and δ_{EW} , obtained by factorizing the lowest order cross section $\hat{\sigma}_0$.

In the partonic cross section, QCD and EW corrections have been factorized. This ansatz is valid up to subleading higher order corrections which start at the 3-loop level (i.e. $\mathcal{O}(\alpha\alpha_s)$ with respect to the lowest order). The factorization of the QCD initial state collinear divergences holds for the hard process described by the electroweak NLO corrections, following from general arguments of the factorization theorems and from the universal nature of the initial state collinear radiation. In fact, the whole set of EW corrections is characterized by a scale m_W , much harder than the one typical of the leading collinear emission. In addition, in the limit of light Higgs, the EW corrections can be expanded as a Taylor series in powers of m_H/m_W and the EW corrections vertex becomes effectively pointlike. In this regime the factorization of the QCD collinear divergences becomes rigorous. For heavier Higgs masses, the factorization should still be valid only in leading order, due to the modifications induced by the EW form factor.

3 Numerical Results

The hadronic proton-proton cross section has been calculated at LHC energy, i.e. $\sqrt{S} = 14$ TeV, in NNLO-QCD accuracy, i.e. setting $\delta_{EW} = 0$, using the MRST2002 NNLO parton distribution functions [13]. The theoretical uncertainty due to the choice of the renormalization scale μ and of the factorization scale M has been canonically estimated by setting $M = \mu$ equal to $m_H/2, m_H, 2m_H$ respectively. The predictions, shown in Fig. 2 (dotted lines), vary by approximately $\pm 8\%$ with respect to the central value. This uncertainty is further reduced when including the effect of the resummation of all the initial state soft gluon radiation [7].

The two-loop electroweak corrections have been added according to Eq. (1) and setting $M = \mu = m_H$. The light fermion corrections can be evaluated for any choice of m_H , whereas the top quark contribution has been computed by means of a Taylor expansion and is limited to the region $m_H \leq 160$ GeV. The hadronic cross section increases from 4 to 8%, for $m_H \leq 160$ GeV. As we can observe in Fig. 2, the effect of the electroweak corrections is an increase of the cross section by an amount which is of the same order of magnitude of the NNLO-QCD theoretical uncertainty, and possibly larger than the uncertainty estimated after the resummation of soft gluon radiation. The main source of uncertainty on the hadronic cross section remains in the accurate determination of the parton distribution functions of the proton.

The effect of the NLO-EW corrections is of great interest, because it enhances the most important Higgs production mechanism and, in turn, affects the absolute number of events of all the Higgs decay modes.

Following Eq. (1), the NLO-EW corrections can be implemented as a simple rescaling of the QCD hadronic cross section. This multiplicative factor is presented in Table 3 as a function of m_H and can be fitted, in the range $114 \text{ GeV} \lesssim m_H \lesssim 155 \text{ GeV}$, by the following simple formula:

$$\delta_{EW}(m_H) = 0.00961 + 6.9904 \cdot 10^{-5} m_H + 2.31508 \cdot 10^{-6} m_H^2. \quad (2)$$

The computation of the NLO-EW corrections to the gluon fusion process has been described in detail in [11, 12]. The analytical expression of the probability amplitude has




Figure 2: The cross section $\sigma_{p+p \rightarrow H+X}$, in pb, is plotted as a function of the mass of the Higgs boson, between 114 GeV and 300 GeV. The dotted lines describe the band of NNLO-QCD uncertainty, for three values of the QCD factorization/renormalization scale $\mu = m_H/2, m_H, 2m_H$. The solid line is the NNLO-QCD ($\mu = m_H$) with the two-loop EW corrections, according to Eq. (1). The two-loop EW corrections include also the top-quark effect, for $m_H \leq 155$ GeV, but only the light quarks contribution for larger values of m_H .

m_H (GeV)	δ_{EW}	m_H (GeV)	δ_{EW}	m_H (GeV)	δ_{EW}	m_H (GeV)	δ_{EW}
114	0.048	136	0.062	158	0.077	180	0.020
116	0.049	138	0.063	160	0.069	182	0.010
118	0.050	140	0.065	162	0.063	184	0.010
120	0.051	142	0.066	164	0.049	186	0.002
122	0.053	144	0.068	166	0.041	188	0.997
124	0.054	146	0.069	168	0.035	190	0.994
126	0.055	148	0.071	170	0.031	192	0.991
128	0.056	150	0.073	172	0.028	194	0.989
130	0.058	152	0.074	174	0.026	196	0.987
132	0.059	154	0.076	176	0.024	198	0.986
134	0.060	156	0.077	178	0.022	200	0.985

Table 1: Rescaling factor δ_{EW} as a function of the Higgs boson mass.

been expressed in terms of Generalized Harmonic PolyLogarithms (GHPL) [14] and has been implemented in a FORTRAN routine¹. The GHPL can be evaluated numerically in several different ways: by direct numerical integration of the basic functions, by power expansions or by solving the associated differential equations. We have checked that these fully independent approaches agree.

4 Conclusions

In conclusion, the calculation of the QCD corrections to the production of a Higgs boson via gluon-fusion has reached a very high level of accuracy; the inclusion of the two-loop electroweak corrections, whose typical size for $m_H \leq 160$ GeV is larger than 5% and then comparable or larger than the QCD uncertainty, is highly desirable. The main source of uncertainty on the hadronic cross section remains in the accurate determination of the parton distribution functions of the proton.

Acknowledgments

The authors wish to thank S. Catani, D. de Florian and M. Grazzini for allowing the use of the numerical program of JHEP **0105** (2001) 025 [arXiv:hep-ph/0102227], and for useful discussions.

This work was partly supported by the European Union under the contract HPRN-CT2002-00311 (EURIDICE) and by MCYT (Spain) under Grant FPA2004-00996, by Generalitat Valenciana (Grants GRUPOS03/013 and GV05/015).

¹available upon request from the authors

References

- [1] H. M. Georgi, S. L. Glashow, M. E. Machacek and D. V. Nanopoulos, Phys. Rev. Lett. **40** (1978) 692.
- [2] S. Dawson, Nucl. Phys. B **359** (1991) 283.
A. Djouadi, M. Spira and P. M. Zerwas, Phys. Lett. B **264** (1991) 440.
- [3] M. Spira, A. Djouadi, D. Graudenz and P. M. Zerwas, Nucl. Phys. B **453** (1995) 17 [arXiv:hep-ph/9504378].
- [4] M. Kramer, E. Laenen and M. Spira, Nucl. Phys. B **511**, 523 (1998) [arXiv:hep-ph/9611272].
- [5] A. Djouadi and P. Gambino, Phys. Rev. Lett. **73** (1994) 2528 [arXiv:hep-ph/9406432].
A. Djouadi, P. Gambino and B. A. Kniehl, Nucl. Phys. B **523** (1998) 17 [arXiv:hep-ph/9712330].
- [6] R. V. Harlander, Phys. Lett. B **492** (2000) 74. [arXiv:hep-ph/0007289];
S. Catani, D. de Florian and M. Grazzini, JHEP **0105** (2001) 025 [arXiv:hep-ph/0102227];
R. V. Harlander and W. B. Kilgore, Phys. Rev. D **64** (2001) 013015 [arXiv:hep-ph/0102241];
R. V. Harlander and W. B. Kilgore, Phys. Rev. Lett. **88** (2002) 201801 [arXiv:hep-ph/0201206];
C. Anastasiou and K. Melnikov, Nucl. Phys. B **646** (2002) 220 [arXiv:hep-ph/0207004];
V. Ravindran, J. Smith and W. L. van Neerven, Nucl. Phys. B **665** (2003) 325 [arXiv:hep-ph/0302135].
- [7] S. Catani, D. de Florian, M. Grazzini and P. Nason, JHEP **0307** (2003) 028 [arXiv:hep-ph/0306211].
- [8] C. Anastasiou, L. J. Dixon and K. Melnikov, Nucl. Phys. Proc. Suppl. **116**, 193 (2003) [arXiv:hep-ph/0211141].
C. Anastasiou, K. Melnikov and F. Petriello, Phys. Rev. Lett. **93** (2004) 262002 [arXiv:hep-ph/0409088].
- [9] R. K. Ellis, I. Hinchliffe, M. Soldate and J. J. van der Bij, Nucl. Phys. B **297** (1988) 221.
S. Catani, E. D’Emilio and L. Trentadue, Phys. Lett. B **211**, 335 (1988).
I. Hinchliffe and S. F. Novaes, Phys. Rev. D **38**, 3475 (1988).
U. Baur and E. W. N. Glover, Nucl. Phys. B **339** (1990) 38.
R. P. Kauffman, Phys. Rev. D **44**, 1415 (1991); Phys. Rev. D **45**, 1512 (1992).
C. P. Yuan, Phys. Lett. B **283**, 395 (1992).
D. de Florian, M. Grazzini and Z. Kunszt, Phys. Rev. Lett. **82** (1999) 5209 [arXiv:hep-ph/9902483].
C. Balazs and C. P. Yuan, Phys. Lett. B **478**, 192 (2000) [arXiv:hep-ph/0001103].
C. Balazs, J. Huston and I. Puljak, Phys. Rev. D **63**, 014021 (2001)

- [arXiv:hep-ph/0002032].
D. de Florian and M. Grazzini, Phys. Rev. Lett. **85**, 4678 (2000) [arXiv:hep-ph/0008152]; Nucl. Phys. B **616**, 247 (2001) [arXiv:hep-ph/0108273].
V. Ravindran, J. Smith and W. L. Van Neerven, Nucl. Phys. B **634** (2002) 247 [arXiv:hep-ph/0201114].
C. J. Glosser and C. R. Schmidt, JHEP **0212** (2002) 016 [arXiv:hep-ph/0209248].
E. L. Berger and J. w. Qiu, Phys. Rev. D **67**, 034026 (2003) [arXiv:hep-ph/0210135]; Phys. Rev. Lett. **91**, 222003 (2003) [arXiv:hep-ph/0304267].
A. Kulesza and W. J. Stirling, JHEP **0312**, 056 (2003) [arXiv:hep-ph/0307208].
A. Kulesza, G. Sterman and W. Vogelsang, Phys. Rev. D **69**, 014012 (2004) [arXiv:hep-ph/0309264].
A. Gawron and J. Kwiecinski, Phys. Rev. D **70**, 014003 (2004) [arXiv:hep-ph/0309303].
G. Watt, A. D. Martin and M. G. Ryskin, Phys. Rev. D **70**, 014012 (2004) [Erratum-ibid. D **70**, 079902 (2004)] [arXiv:hep-ph/0309096].
J. Smith and W. L. van Neerven, Nucl. Phys. B **720**, 182 (2005) [arXiv:hep-ph/0501098].
C. Anastasiou, K. Melnikov and F. Petriello, Nucl. Phys. B **724** (2005) 197 [arXiv:hep-ph/0501130].
A. V. Lipatov and N. P. Zotov, Eur. Phys. J. C **44**, 559 (2005) [arXiv:hep-ph/0501172].
- [10] G. Bozzi, S. Catani, D. de Florian and M. Grazzini, Phys. Lett. B **564** (2003) 65 [arXiv:hep-ph/0302104]; Nucl. Phys. B **737** (2006) 73 [arXiv:hep-ph/0508068].
- [11] U. Aglietti, R. Bonciani, G. Degrossi and A. Vicini, Phys. Lett. B **595** (2004) 423 [arXiv:hep-ph/0404071]; Phys. Lett. B **600** (2004) 57 [arXiv:hep-ph/0407162].
- [12] G. Degrossi and F. Maltoni, Phys. Lett. B **600** (2004) 255 [arXiv:hep-ph/0407249].
- [13] A. D. Martin, R. G. Roberts, W. J. Stirling and R. S. Thorne, Phys. Lett. B **531** (2002) 216 [arXiv:hep-ph/0201127].
- [14] U. Aglietti and R. Bonciani, Nucl. Phys. B **668** (2003) 3 [arXiv:hep-ph/0304028]; Nucl. Phys. B **698** (2004) 277 [arXiv:hep-ph/0401193].